

Magnetic order in the Ising model with parallel dynamics

Emilio N.M. Cirillo

Dipartimento Me. Mo. Mat., Università degli Studi di Roma “La Sapienza,” via A. Scarpa 16, 00161 Roma, Italy

Francesca R. Nardi

Eurandom, PO BOX 513, 5600MB, Eindhoven, Netherlands

Antonio D. Polosa

Department of Physics, University of Helsinki, PO BOX 64, FIN-00014, Finland

(February 1, 2008)

It is discussed how the equilibrium properties of the Ising model are described by an Hamiltonian with an antiferromagnetic low temperature behavior if only an heat bath dynamics, with the characteristics of a Probabilistic Cellular Automaton, is assumed to determine the temporal evolution of the system.

PACS numbers: 05.50.+q; 75.10.-b; 64.60.Fr

I. INTRODUCTION

In this note we discuss the equilibrium properties of Probabilistic Cellular Automata (PCA) reversible with respect to a Gibbs measure derived by a suitable Hamiltonian. A PCA [1–3] is a lattice model with discrete variables which are subject to a probabilistic *simultaneous* updating in discrete time steps: all configurations are accessible in a single updating.

PCA arise as an extended definition of Deterministic Cellular Automata in which the updating follows a set of deterministic local rules. The huge number of possible deterministic (or probabilistic) rules makes the topic of Cellular Automata overwhelmingly abundant. One of the most famous Cellular Automata systems is the Conway’s “Game of life” [4]; in spite of the very simple deterministic majority rule assigned, the system, which is a kind of spin lattice, presents an extremely rich and complex evolution pattern.

PCA have been studied in a wide variety of contexts, ranging from biology to the theory of automation, while their role in statistical mechanics has been thoroughly investigated only relatively recently in [5–9]. The particular family of automata we study is obtained by implementing in parallel fashion the heat bath dynamics [6]. In other terms, we define a rule for the transition probabilities such that all single spins of a lattice are updated simultaneously with heat bath rates. This amounts to define a Markov chain for the evolution of the spin system, having the characteristics of a PCA.

We observe that the way of implementing the heat bath dynamics reflects into a qualitative modification of the equilibrium properties of the model. In particular, an Ising-like ferromagnetic Hamiltonian with two body interactions, defines a PCA reversible with respect to a Gibbs measure determined by an Hamiltonian allowing a low temperature antiferromagnetic phase. This behavior is absent if a serial dynamics is implemented, for which at most one spin of the system is updated at any time.

The paper is organized as follows. In Section II we define the PCA under consideration and the spin model Hamiltonian, H , determining the heat bath single spin rates. In Section III the structure and the low temperature antiferromagnetic properties of H' are discussed. There follows a discussion of its equilibrium properties and ground states. Finally, in Section IV, it is considered the case in which the heat bath spin rates are determined by a (two body) next-to-nearest neighbor interaction. The dynamical generation of antiferromagnetic couplings is reviewed in the conclusive remarks.

II. COUPLING PROLIFERATION IN REVERSIBLE PCA’S

Let Λ be a finite two-dimensional square lattice and $|\Lambda|$ its cardinality. For each $x = (x_1, x_2), y = (y_1, y_2) \in \Lambda$ we denote by $|x - y|$ the Euclidean distance on the lattice. Let $\sigma(x) \in \{-1, +1\}$ a spin variable associated to the site $x \in \Lambda$; the space $\{1, -1\}^\Lambda$ of configurations is denoted by $\mathcal{S}$.

Let us consider a generic Hamiltonian $H : \sigma \in \mathcal{S} \rightarrow H(\sigma) \in \mathbb{R}$ and the corresponding equilibrium Gibbs measure

$$\mu(\sigma) = \frac{e^{-\beta H(\sigma)}}{\sum_{\eta \in \mathcal{S}} e^{-\beta H(\eta)}} \quad , \quad (1)$$

with β the inverse of the temperature. We now define the heat bath single spin rates: given the site $x \in \Lambda$, we consider the Gibbs equilibrium measure for σ_x with respect to a fixed configuration σ on $\Lambda \setminus \{x\}$. Letting $a \in \{-1, +1\}$, we have

$$p_x(a|\sigma) = \frac{\exp\{-\beta H(a, \sigma)\}}{\exp\{-\beta H(a, \sigma)\} + \exp\{-\beta H(-a, \sigma)\}} \quad , \quad (2)$$

where $(\pm a, \sigma)$ are the configurations equal to σ on $\Lambda \setminus \{x\}$ and to $\pm a$ on x . Note that the normalization condition $p_x(a|\sigma) + p_x(-a|\sigma) = 1$ is trivially satisfied.

We can now implement the heat bath dynamics in a serial fashion, namely we can consider the Markov chain σ_t (t being the discrete time temporal variable), with transition probabilities

$$P(\sigma, \eta) = \begin{cases} (1/|\Lambda|)p_x(\eta(x)|\sigma) & \text{if } \exists x \in \Lambda \text{ such that } \sigma = \eta \text{ on } \Lambda \setminus \{x\} \\ 0 & \text{otherwise} \end{cases} \quad (3)$$

for all $\sigma, \eta \in \mathcal{S}$. The transition probabilities (3) are reversible with respect to the Gibbs measure (1), i.e., the detailed balance condition is satisfied or, equivalently, the equilibrium measure is the Gibbs measure (1).

A different point of view can be taken [6]: we define the transition probabilities $P(\sigma, \eta)$ in such a way that all the spins are *simultaneously* and independently updated, in a parallel fashion, with the heat bath rates (2). Thus, instead of (3) we consider the Markov chain σ_t defined by

$$P(\sigma, \eta) = \prod_{x \in \Lambda} p_x(\eta(x)|\sigma) \quad \forall \sigma, \eta \in \mathcal{S} \quad . \quad (4)$$

This amounts to define a Probabilistic Cellular Automata (PCA).

In general the equilibrium properties of the Markov chain (4) are not trivial, for instance it is not obvious that there exists a Gibbs measure such that the detailed balance principle is satisfied.

Let us consider, now, the case of the two body interactions and suppose that the Hamiltonian has the form:

$$H(\sigma) = - \sum_{x, y \in \Lambda} J_{x, y} \sigma(x) \sigma(y) - \sum_{x \in \Lambda} h_x \sigma(x) \quad , \quad (5)$$

where $J_{x, y} \in \mathbb{R}$ are the pair couplings between spins at sites x, y , and $h_x \in \mathbb{R}$ is the external magnetic field acting on the spin at site x . For physical reasons we suppose that the pair couplings are symmetric, namely $J_{x, y} = J_{y, x}$ for all $x, y \in \Lambda$. The heat bath single spin rates are given by

$$p_x(a|\sigma) = \frac{1}{1 + \exp\{-\beta[H(-a, \sigma) - H(a, \sigma)]\}} = \frac{1}{2} [1 + a \tanh \beta S_x(\sigma)] \quad , \quad (6)$$

where

$$S_x(\sigma) = \sum_{y \in \Lambda \setminus \{x\}} J_{x, y} \sigma(y) + h_x \quad , \quad (7)$$

for any $\sigma \in \mathcal{S}$ and $x \in \Lambda$. It is easy to show [6] that the Probabilistic Cellular Automaton (4) with single spin rates (6) is reversible with respect to the Gibbs measure μ' on $\mathcal{S}$ associated to the Hamiltonian

$$H'(\sigma) = -\beta \sum_{x \in \Lambda} h_x \sigma(x) - \sum_{x \in \Lambda} \log \cosh [\beta S_x(\sigma)] \quad . \quad (8)$$

In other words the detailed balance condition

$$P(\sigma, \eta) \exp\{-H'(\sigma)\} = P(\eta, \sigma) \exp\{-H'(\eta)\} \quad , \quad (9)$$

is satisfied for any $\sigma, \eta \in \mathcal{S}$. This means that $H'(\sigma)$ is the equilibrium Hamiltonian of a system governed by $H(\sigma)$ and evolving with the law (4).

The choice of the two body interaction in equation (5) is strictly connected to the reversibility of the resulting Probabilistic Cellular Automaton (4) [2,3]. For example, consider the three body interaction Hamiltonian $H(\sigma) = \sum_{x,y,z \in \Lambda} J_{x,y,z} \sigma(x)\sigma(y)\sigma(z)$ with the three body couplings $J_{x,y,z}$ symmetric with respect to permutations of the indices and such that $J_{x,y,z} \neq 0$ if and only if $x \neq y \neq z \neq x$. Then the problem of showing the reversibility of the parallel dynamics can be reduced to the problem of finding a function $\phi : \sigma \in \mathcal{S} \rightarrow \phi(\sigma) \in \mathbb{R}$ such that

$$\phi(\sigma) - 3\beta\eta(x) \sum_{y,z \in \Lambda \setminus \{x\}} J_{x,y,z} \sigma(y)\sigma(z) = \phi(\eta) - 3\beta\sigma(x) \sum_{y,z \in \Lambda \setminus \{x\}} J_{x,y,z} \eta(y)\eta(z) \quad ,$$

which has no solution.

Let us now discuss the main feature of reversible heat bath derived probabilistic automata. As it has been seen above, if the starting Hamiltonian is given by (5), then the Markov chain (4) is reversible with respect to the Gibbs measure with Hamiltonian H' given by (8). It is clear that new kind of interactions, different from the one present in the original Hamiltonian H , arise when H' is considered.

Suppose, for instance, that the starting Hamiltonian has range $r > 0$, namely $J_{x,y} = 0$ for any $x, y \in \Lambda$ such that $|x - y| > r$. Then we have $S_x(\sigma) = \sum_{y: 0 < |x-y| \leq r} J_{x,y} \sigma(y) + h_x$, a sort of average of the spins inside a ball centered at site x with radius equal to r . Hence, by expanding H' as a sum of potentials we will get all the possible couplings inside the ball, starting from the two body up to the $N(r)$ body interaction, with $N(r)$ the number of sites inside the ball. In some sense these new couplings are dynamically generated. In the following we will discuss few interesting particular cases.

III. NEAREST NEIGHBOR ISING MODEL

Let us consider the standard nearest neighbors Ising model with no external magnetic field, namely we consider (5) with $J_{x,y} = J/2$ for any $x, y \in \Lambda$ such that $|x - y| = 1$, $J_{x,y} = 0$ otherwise, and $h_x = 0$ for any $x \in \Lambda$. The Hamiltonian H' is the sum of averages performed over the four site crosses centered at each site of the lattice. We then expect all the possible interactions inside the cross.

As it has been seen in [10] it is possible to extract the potentials and rewrite the Hamiltonian H' in the following way

$$H'(\sigma) = -J_1 \sum_{\langle xy \rangle_{\sqrt{2}}} \sigma(x)\sigma(y) - J_2 \sum_{\langle xy \rangle_2} \sigma(x)\sigma(y) - J_3 \sum_{\diamond_{xywz}} \sigma(x)\sigma(y)\sigma(w)\sigma(z) \quad , \quad (10)$$

where the three sums (see Fig. 1a) are respectively performed over the pairs of next to the nearest neighbors (sites at distance $\sqrt{2}$), the pairs of third neighbors (sites at distance 2), the four site diamond shaped clusters (plaquettes with side length equal to $\sqrt{2}$). The coupling constants are given by

$$J_1 = \frac{1}{4} \log \cosh(2\beta J) \sim \frac{1}{2}\beta J, \quad J_2 = \frac{1}{2}J_1, \quad J_3 = \frac{1}{16} \log \frac{\cosh^2(2\beta J)}{\cosh^8(\beta J)} \sim -\frac{1}{4}\beta J \quad , \quad (11)$$

where “ $\sim$ ” means the limiting behavior for $\beta \rightarrow \infty$. There exist several possible ways to extract the potentials. A very natural one, in the case of spin variable, is to consider the function $\varphi_x(\sigma) = \log \cosh[(\beta J/2) \sum_{y \in \Delta_x} \sigma(y)]$, where $\Delta_x = \{y \in \Lambda : |y - x| = 1\}$ is the set of nearest neighbors of site x , and its expansion

$$\varphi_x(\sigma) = \sum_{X \subset \Delta_x} c(X) \prod_{y \in X} \sigma(y) \quad ,$$

with the coefficients $c(X)$ given by

$$c(X) = \frac{1}{2^{|\Delta_x|}} \sum_{\sigma \in \{-1, +1\}^{\Delta_x}} \varphi_x(\sigma) \prod_{y \in X} \sigma(y) \quad ,$$

where, we recall, $|\Delta_x| = 4$ is the cardinality of Δ_x (see [11]).

It is important to remark that the second nearest neighbor interaction, J_1 , is positive and dominating; hence we expect a low temperature antiferromagnetic phase to exist. What appears very remarkable is that we have derived an antiferromagnetic behavior in a purely dynamical way as a result of the coupling proliferation. If a parallel heat

bath Ising dynamics is implemented, the equilibrium Gibbs measure shows a low temperature antiferromagnetic phase despite the simple physical ferromagnetic coupling of the Ising model. This phenomenon is, obviously, absent if the Ising heat bath dynamics is implemented in a serial fashion.

The equilibrium properties of the model can be understood by remarking that two independent models are found if the lattice is partitioned into two square sublattices with step $\sqrt{2}$ (the even and the odd sublattice). Each model is, indeed, an eight vertex model with nearest neighbors coupling J_1 , next to the nearest neighbors coupling J_2 and plaquette interaction J_3 . This model has been widely studied both in two [12–14] and three [15,16] dimensions. From (11) and the very well known properties of the two-dimensional eight vertex model we have that on each sublattice there are two coexisting low temperature phases, respectively with positive and negative magnetization. Hence, by combining in all the possible ways the two phases we get, for the original model, the 3 different low temperature phases corresponding to the three ground states ψ_0, ψ_1, ψ_2 (see Fig. 2).

It is of some interest a direct study of the hamiltonian (10): ground states can be defined as those configurations on which the Gibbs measure μ' , associated to the Hamiltonian H' , is concentrated when the limit $\beta \rightarrow \infty$ is considered, namely as the minima of the energy

$$E(\sigma) = \lim_{\beta \rightarrow \infty} \frac{H'(\sigma)}{\beta} = - \sum_{x \in \Lambda} |S_x(\sigma)| \quad , \quad (12)$$

uniformly in $\sigma \in \mathcal{S}$. It is rather clear that with periodic boundary conditions there exist three coexisting minima $\psi_0, \psi_1, \psi_2 \in \mathcal{S}$ (see Fig. 2), with energy $-4|\Lambda|$, such that $\psi_0(x) = +1$, $\psi_1(x) = (-1)^{x_1+x_2}$ and $\psi_2(x) = -1$ for all $x = (x_1, x_2) \in \Lambda$. Notice that ψ_1 is the chessboard configuration.

The problem is, now, to understand if this coexistence of different states persists at a finite small temperature, namely if the system undergoes a low temperature phase transition. We give an heuristic argument: at finite temperature ground states are perturbed because small droplets of different phases show up. The idea is to calculate the energetic cost of a perturbation of one of the four coexisting states via the formation of a square droplet of a different phase. If it results that one of the three states ψ_0, ψ_1, ψ_2 is more easily perturbed, then we will conclude that this is the equilibrium phase at finite temperature.

A simple calculation, see [10], shows that the energy cost of a square droplet of side length n of one of the two homogeneous ground states plunged in one of the two chessboards (or vice versa) is equal to $8n$. On the other hand if an homogeneous phase is perturbed as above by the other homogeneous phase, or one of the two chessboards is perturbed by the other one, then the energy cost is $16n$.

Hence, from the energetical point of view the most convenient excitations are those in which an homogeneous phase is perturbed by a chessboard or vice versa. Moreover, for each state ψ_0, ψ_1, ψ_2 there exist two possible energetically convenient excitations: there is no entropic reason to prefer one of the four ground states to the others when a finite low temperature is considered. This remark strongly suggests that at small finite temperature the three ground states still coexist.

IV. NEXT TO THE NEAREST NEIGHBOR ISING MODEL

Let us consider the Ising model with no external magnetic field and next to the nearest neighbor interaction, namely we consider (5) with $J_{x,y} = J/2$ for any $x, y \in \Lambda$ such that $|x - y| = \sqrt{2}$, $J_{x,y} = 0$ otherwise, and $h_x = 0$ for any $x \in \Lambda$.

It is possible to extract the potentials as seen in Section III. The Hamiltonian H' can be written as

$$H'(\sigma) = -J_1 \sum_{\langle xy \rangle_2} \sigma(x)\sigma(y) - J_2 \sum_{\langle xy \rangle_{2\sqrt{2}}} \sigma(x)\sigma(y) - J_3 \sum_{\square_{xyzw}} \sigma(x)\sigma(y)\sigma(w)\sigma(z) \quad , \quad (13)$$

where the three sums (see Fig. 1b) are respectively performed over the pairs of third nearest neighbors (sites at distance 2), the pairs of sites at distance $2\sqrt{2}$, the plaquettes with side length equal to 2. The coupling constants are still given by (11).

In order to study this model we remark that if the lattice is partitioned into four square sublattices with step 2, then we obtain four independent models one on each sublattice, each model being again an eight vertex model with nearest neighbors coupling J_1 , next to the nearest neighbors coupling J_2 and plaquette interaction J_3 . Hence, on each sublattice we have the two degenerate ground states with all the spins, respectively, equal to one and minus one. By combining in all the possible ways these two states we get, for our model, $2^4 = 16$ different ground states. On the torus, namely when periodic boundary conditions are considered, some of the ground states are equivalent, so we get

the seven states $\psi_0, \psi_1, \dots, \psi_6$ in Fig. 2 and 3. The fact that the phase transition persists at finite small temperature is, as in Section III a straightforward consequence of the known behavior of the eight vertex model.

In this note we observed the relation between the two body ferromagnetic interaction of the Ising model and the low temperature antiferromagnetic behavior of the equilibrium Hamiltonian obtained evolving the initial system with a parallel heat bath dynamics. Any lattice system with two body interactions and having a self-organization resembling the parallel dynamics here described in the simple Ising model, could evolve towards unexpectedly complex equilibrium states. The influence of the parallel dynamics on the physical properties of the equilibrium measure emerges as an interesting new feature that deserves further explorations.

ACKNOWLEDGMENTS

ENMC wishes to express his thanks to the Physics Department of the University of Helsinki for its warm hospitality. ENMC also thanks J.L. Lebowitz for having introduced him to the interesting topic of PCA's and P. Dai Pra for a useful discussion. ADP acknowledges support from EU-TMR programme, contract CT98-0169.

-
- [1] O.N. Stavskaja, *Math. USSR Sobornik* **21**, 395–411 (1973).
 - [2] V. Kozlov, “Reversible Markov chain with local interactions,” in “Multicomponent random system,” Adv. in Prob. and Rel. Topics, 1980.
 - [3] Vasiljev, “Bernoulli and Markov stationary measures in discrete local interactions,” Lect. Notes in Math. 653, 1978.
 - [4] M. Creutz, hep-lat/9611017
 - [5] P. Rujan, *J. Stat. Phys.* **49**, 139–222 (1987); J.L. Lebowitz, C. Maes and E. Speer, *J. Stat. Phys.* **59**, 117 (1990).
 - [6] B. Derrida, “Dynamical phase transition in spin model and automata,” Fundamental problem in Statistical Mechanics VII, H. van Beijeren, Editor, Elsier Science Publisher B.V., (1990).
 - [7] J.L. Lebowitz, C. Maes, E. Speer, *J. Stat. Phys.* **59**, 117–170 (1990).
 - [8] J.L. Lebowitz, C. Maes, E. Speer, “Probabilistic cellular automata: some statistical mechanics considerations,” in *Lectures in Complex Systems, SFI Studies in the Sciences of Complexity, Lecture Volume II*, ed. E. Jen (Addison Wesley, New York, 1990).
 - [9] S. Bigelis, E.N.M. Cirillo, J.L. Lebowitz, E.R. Speer, *Phys. Rev. E* **59**, 3935 (1999).
 - [10] E.N.M. Cirillo, F.R. Nardi, “Metastability in the Ising model with parallel dynamics,” Preprint 2001.
 - [11] K. Haller, T. Kennedy, *J. Stat. Phys.* **85**, 607–637 (1996).
 - [12] R.J. Baxter, *Phys. Rev. Lett.* **26**, 832 (1971).
 - [13] R.J. Baxter, “Exactly solved models in Statistical Mechanics,” (Academic Press, London, 1982).
 - [14] M. Nauenberg, B. Nienhuis, *Phys. Rev. Lett.* **33**, 944 (1974).
 - [15] A. Cappi, P. Colangelo, G. Gonnella, A. Maritan, *Nucl. Phys. B* **370**, 659 (1992).
 - [16] E.N.M. Cirillo, G. Gonnella, *Journ. Phys. A: Math. Gen.* **28**, 867 (1995).

Figure captions

Fig. 1: Couplings J_1 , J_2 and J_3 for the hamiltonian (10) (resp. (13)) are shown in (a) (resp. (b)).

Fig. 2: The three ground states ψ_0, ψ_1, ψ_2 depicted from the left to the right.

Fig. 3: The four ground states $\psi_3, \psi_4, \psi_5, \psi_6$ depicted from the left to the right.




Fig. 1




Fig. 2




Fig. 3